



Original Article

Asian Pacific Journal of Tropical Medicine

apjtm.org



doi:10.4103/1995–7645.318304

Impact Factor: 1.94

Ivermectin as an adjunct treatment for hospitalized adult COVID–19 patients: A randomized multi–center clinical trial

Morteza Shakhshi Niaee¹, Peyman Namdar², Abbas Allami³, Leila Zolghadr⁴, Amir Javadi⁵, Amin Karampour³, Mehran Varnaseri⁶, Behzad Bijani⁷, Fatemeh Cheraghi¹, Yazdan Naderi⁸, Fatemeh Amini⁹, Masoumeh Karamyan⁶, Mohammad Jafar YadYad⁶, Ramin Jamshidian⁶, Nematollah Gheibi^{8✉}

¹Sciences and Technology Park, Qazvin, Iran

²Department of Surgery, Qazvin University of Medical Sciences, Qazvin, Iran

³Department of Infectious Diseases, Clinical Research Development Unit, BouAliSina Hospital, Qazvin University of Medical Sciences, Qazvin, Iran

⁴Department of Chemistry, Imam Khomeini International University Qazvin, Iran

⁵Department of Social Sciences School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

⁶Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁷Department of Infectious Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

⁸Cellular and Molecular Research Center, Research Institute for Prevention of Non–Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

⁹Department of Persian Medicine, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

ABSTRACT

Objective: To evaluate different doses of ivermectin in adult patients with mild COVID-19 and to evaluate the effect of ivermectin on mortality and clinical consequences.

Methods: A randomized, double-blind, placebo-controlled, multicenter clinical trial was performed at five hospitals. A total of 180 mild hospitalized patients with COVID-19 confirmed by PCR or chest image tests were enrolled and allocated to six arms including hydroxychloroquine 200 mg twice per day, placebo plus hydroxychloroquine 200 mg twice per day, single dose ivermectin (200 µg/kg), three low interval doses of ivermectin (200, 200, 200 µg/kg), single dose ivermectin (400 µg/kg), and three high interval doses of ivermectin (400, 200, 200 µg/kg). The primary endpoint of this trial was all-cause of mortality or clinical recovery. The radiographic findings, hospitalization and low O₂ saturation duration, and hematological variables of blood samples were analyzed.

Results: A total of 16.7% (5/30) and 20.0% (6/30) patients died in arms treated with hydroxychloroquine 200 mg twice per day and placebo plus hydroxychloroquine 200 mg twice per day, respectively, and a reduction in mortality rate in patients receiving ivermectin treatment to 0%, 10%, 0% and 3.3% for arms 1–4 were observed. Risk of mortality was also decreased about 15% in the ivermectin treated arms.

Conclusions: Ivermectin as an adjunct reduces the rate of mortality, time of low O₂ saturation, and duration of hospitalization in adult COVID-19 patients. The improvement of other clinical parameters

shows that ivermectin, with a wide margin of safety, had a high therapeutic effect on COVID-19.

KEYWORDS: SARS-COV-2; Ivermectin; Randomized controlled trial; Dose-respond relationship

1. Introduction

The COVID-19 disease has become a pandemic after the WHO declaration in March 2020. This disease has created a difficult condition around the world, and hence there is an important and urgent need to find proper treatments for an effective cure, decrease the virus carriage duration, and thus limit its transmission in society[1–3]. So far, different drugs such as hydroxychloroquine, azithromycin, remdesivir, oseltamivir, lopinavir, and ritonavir have been used against COVID-19[4–10]. However, among the candidate treatments, only remdesivir have been tested in large

✉To whom correspondence may be addressed. E-mail: gheibi_n@yahoo.com, ngheibi@qums.ac.ir

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2021 Asian Pacific Journal of Tropical Medicine Produced by Wolters Kluwer-Medknow. All rights reserved.

How to cite this article: Shakhshi Niaee M, Namdar P, Allami A, Zolghadr L, Javadi A, Karampour A, et al. Ivermectin as an adjunct treatment for hospitalized adult COVID-19 patients: A randomized multi-center clinical trial. Asian Pac J Trop Med 2021; 14(6): 266–273.

Article history: Received 12 December 2020
Accepted 4 June 2021

Revision 31 May 2021
Available online 25 June 2021